

Surgico-Pediatric Conference September 12, 2006

Nuthapong UKarapol, MD

Peptic Ulcer disease in children

- Primary PUD – uncommon in childhood
- 2% (52 out of 2550)
 - 10 Gastric ulcer
 - 42 Duodenal ulcer
- Mean age 10.5 yr
- No specific symptoms and signs
- Associated with Hp in 54%

Roma E, et al. Eur J Pediatr 2001;160:497.

Alarm symptoms and signs

- Involuntary weight loss
- Deceleration of linear growth
- Gastrointestinal blood loss
- Significant vomiting
- Chronic severe diarrhea
- Persistent right upper or right lower quadrant pain
- Unexplained fever and family history of IBD

Leading points in childhood intussusception

- Found in 31/567 cases – 5.4%
- Average age is twice that of children whose intussusceptions are idiopathic
- Harder to diagnose
- Duration of symptoms and signs is longer than is usually seen
- Barium enema as a treatment modality is usually not successful
- More common in small bowel intussusception (40%)
- Higher morbidity

Ein SH. J Pediatr Surg 1976;11:209

Leading points in childhood intussusception

LEADING PATHOLOGY	N (total 30)
Meckel's diverticulum	14
Polyp	8
Ileum	6
Cecum	1
Colon	1
Duplication	5
Ileum	2
Cecum	2
Jejunum	1
Henoch-Schonlein purpura	1
Suture line	1
Appendix	1
Lymphoma	1

Ein SH. J Pediatr Surg 1976;11:209

Intussusception due to GI lymphoma

- Found in 11/1200 cases – 0.9% (1953–1984)
- Age seems to be as important a clue
- Longer duration of symptoms
- Rarely be reduced by hydrostatic barium enema

Ein SH. J Pediatr Surg 1986;21:786

Intussusception due to GI lymphoma

Intussusception vs lymphoma	
Age, yrs mean (range)	7 (3-13)
Site	
Small bowel intussusception	73%
Colonic intussusception	27%
Duration, days mean (range)	42 (1-180)
Symptoms	
Abdominal pain	91%
Abdominal mass	73%
Mortality rate	73%

Ein SH. J Pediatr Surg 1986;21:786

Intussusception in Adults

- 5% of all intussusception
- 1%-5% of all causes of intestinal obstruction in adults
- Over 90% is associated with precipitating causes
- Clinical presentation is more chronic (57%)
- Palpable mass and rectal bleeding are less likely to be observed (<1/3), compared to children
- Only half (52%) of the cases can be correctly diagnosed preoperatively
- CT scan seems to be the most useful diagnostic tool for this group of patients

Goh BKP, et al. World J Surg 2006;30:1300

Intussusception in Adults

TYPE of intussusception	%
Enteric	51.7%
Jejunojejunal	20%
Ileoileal	6.7%
Ileocolic	2.5%
Colonic	48.3%
Ileocecal-colic	13.3%
Colocolic	23.3%
Sigmoidorectal	6.7%
Appedicocecal	5.0%

Goh BKP, et al. World J Surg 2006;30:1300

Intussusception in Adults - Benign leading points

Small bowel intussusception	Colonic intussusception
Peutz-Jegher polyp	Adenoma
Adenoma	Inflammatory pseudopolyp
Leiomyoma	Lipoma
Crohn's disease	
Lipoma	
Inflammatory pseudopolyp	
Neurofibroma	
Meckel's	
Tuberculosis	
Indeterminate polyp	

Goh BKP, et al. World J Surg 2006;30:1300

Intussusception in Adults - Malignant leading points

Small bowel intussusception	Colonic intussusception
Metastatic lung carcinoma	Adenocarcinoma
Metastatic melanoma	Lymphoma
Metastatic gastric carcinoma	Metastatic melanoma
Adenocarcinoma	Malignant GIST
Lymphoma	
Malignant GIST	

Goh BKP, et al. World J Surg 2006;30:1300

Intussusception in Adults - Predictive factors for malignancy as a leading point

	Benign (n=32)	Malignant (n=28)	P	P
Age, yrs; median	50.2	69.0	0.009	
Sex (M:F)	1:1	1.8:1	0.265	
Duration of symptoms			0.265	
Acute (<14 d)	16	10		
Chronic (>14 d)	16	18		
Rectal bleeding or melena	5 (15.6%)	9 (32.1%)	0.131	
Anemia (Hb <12 g/dl)	6 (18.8%)	20 (71.4%)	<.001	0.001
Site			0.001	0.004
Enteric	23 (71.9%)	8 (28.6%)		
Colon	9 (28.1%)	20 (71.4%)		


Goh BKP, et al. World J Surg 2006;30:1300

Intussusception in Adults

- Advocate en bloc resection in all patients with colonic intussusception because of its high malignancy rate (69%) and furthermore 16 of 20 malignant lesions (80%) are primary malignancies
- Reduction followed by resection may be applied to small bowel lesion, as only 26% of these lesions are malignant and 6 of 8 (75%) are metastatic cancers or lymphoma


Goh BKP, et al. World J Surg 2006;30:1300

GI Non-Hodgkin's lymphoma



Halme L, et al. Acta Oncologica 1997;36:69

GI Non-Hodgkin's lymphoma



Kolve M, et al. Gastrointest Endosc 1999;49:307.

GI Non-Hodgkin's lymphoma

- Helicobacter pylori infection
- HIV (Burkitt's lymphoma)
- Solid organ transplantation
- Celiac disease
- Inflammatory bowel disease (colonic lymphoma)
- HCV infection – controversial

Halme L, et al. Acta Oncologica 1997;36:69

Take Home Message

- Carefully look for alarm symptoms in any patient presenting with chronic abdominal pain, in order to offer proper management
- Anemia, hypoalbuminemia, edema, and chronic abdominal pain support chronic blood loss (protein losing enteropathy) in this patient
- Unidentified intussusception by barium enema cannot exclude small bowel intussusception
- Intussusception in older children and adults need further investigation to find out a possibility of leading points
- Although malignancy as a cause of leading point for intussusception is rare in children, it should be considered, particularly in a patient with anemia.